



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Atypical Femoral Fracture in Osteoporosis Pseudoglioma Syndrome Associated With Two Novel Compound Heterozygous Mutations in LRP5

Citation for published version:

Alonso, N, Soares, DC, McCloskey, E, Summers, GD, Ralston, SH & Gregson, CL 2015, 'Atypical Femoral Fracture in Osteoporosis Pseudoglioma Syndrome Associated With Two Novel Compound Heterozygous Mutations in LRP5', *Journal of Bone and Mineral Research*, vol. 30, no. 4, pp. 615-620.
<https://doi.org/10.1002/jbmr.2403>

Digital Object Identifier (DOI):

[10.1002/jbmr.2403](https://doi.org/10.1002/jbmr.2403)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Bone and Mineral Research

Publisher Rights Statement:

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



**ATYPICAL FEMORAL FRACTURE IN OSTEOPOROSIS PSEUDOGLIOMA SYNDROME
ASSOCIATED WITH TWO NOVEL COMPOUND HETEROZYGOUS MUTATIONS IN *LRP5*[†]**

Nerea Alonso¹, Dinesh C. Soares², Eugene McCloskey³, Gregory D. Summers⁴, Stuart H. Ralston¹, Celia L. Gregson^{5*}

¹ Rheumatic Diseases Unit, Centre for Genomic and Experimental Medicine, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

² MRC Human Genetics Unit and Centre for Genomic and Experimental Medicine, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK

³ Academic Unit of Bone Metabolism, Centre for Biomedical Research, Northern General Hospital, Sheffield, UK

⁴ Department of Rheumatology, Derby Hospitals NHS Foundation Trust, Derby, UK

⁵ Musculoskeletal Research Unit, School of Clinical Sciences, University of Bristol, Bristol, UK

* Correspondence to: Dr Celia L. Gregson

Musculoskeletal Research Unit, School of Clinical Sciences, University of Bristol, Learning & Research Building (Level 1), Bristol, BS10 5NB, United Kingdom

Tel: +44(0) 117 4147842

Fax: +44(0) 117 4147924

Email: celia.gregson@bristol.ac.uk

DISCLOSURE

Authors have no conflicts of interest to declare.

[†]This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/jbmr.2403]

Additional Supporting Information may be found in the online version of this article.

Initial Date Submitted August 15, 2014; Date Revision Submitted October 20, 2014; Date Final Disposition Set November 5, 2014

Journal of Bone and Mineral Research

© 2014 American Society for Bone and Mineral Research

DOI 10.1002/jbmr.2403

ABSTRACT

Osteoporosis pseudoglioma syndrome (OPPG) is a rare autosomal recessive condition of congenital blindness and severe childhood osteoporosis with skeletal fragility, caused by loss-of-function mutations in the *LRP5* (Low-density lipoprotein receptor-related protein 5) gene. We report the first case of atypical (subtrochanteric) femoral fracture (AFF) in OPPG, occurring in a 38 year old man within the context of relatively low bone turnover and trabecular osteoporosis on bone histology. We identify two novel *LRP5* mutations: *R752W* is associated with low bone mass density (BMD), as demonstrated by the heterozygous carriage identified in his 57 year old mother; however, the combination of this *R752W* mutation with another novel *W79R* mutation, causes a severe case of compound heterozygous OPPG. We undertake three-dimensional homology modelling of the four extracellular YWTD β -propeller/EGF-like domains (E1-E4) of LRP5, and show that both novel mutations destabilise the β -propeller domains that are critical for protein and ligand binding to regulate Wnt signalling and osteoblast function. Whilst AFFs have been reported in other rare bone diseases, this is the first in a genetic condition of primary osteoblast dysfunction. The relatively low bone turnover observed, and knowledge of *LRP5* function, implicates impaired bone remodelling in the pathogenesis of AFF.

KEY WORDS

Atypical femoral fracture, compound heterozygote, LRP5, OPPG, protein model

INTRODUCTION

Atypical femoral fractures (AFF) have been reported in patients taking anti-resorptive therapy, particularly for prolonged durations, although they can occur in bisphosphonate-naïve individuals. The pathogenesis of AFF remains unclear and much anxiety exists regarding the association between commonly used osteoporosis treatments and AFF⁽¹⁾. Studying rare genetic diseases, such as *LRP5* disorders, can offer insights into the understanding of more common diseases.

Loss-of-function *LRP5* mutations cause osteoporosis pseudoglioma syndrome⁽²⁾ (OPPG, OMIM:259770), a rare autosomal recessive disorder, characterised by congenital or juvenile-onset blindness, severe juvenile-onset osteoporosis^(3;4), skeletal fragility⁽⁵⁾ and occasionally learning difficulties⁽⁶⁾. Approximately 70 cases have been reported worldwide⁽⁶⁻¹⁰⁾. Ocular abnormalities are detected at birth or in the first few years of life. Short stature and bone deformities can develop. Biochemical markers of bone resorption and formation are usually normal^(5;11). Heterozygous carriers often have decreased bone mass but normal eye sight^(2;12;13). Activating *LRP5* mutations confer the opposing phenotype of high bone mass and fracture resistance (HBM, OMIM:601884)⁽¹⁴⁻¹⁷⁾.

LRP5 is a cell-surface receptor which activates the canonical Wnt/ β -catenin pathway, regulating osteoblastic bone formation⁽¹⁰⁾. The *LRP5* protein carries an extracellular domain composed of four β -propellers and EGF motifs and three LDL-receptor domains, a 23-amino acid membrane-spanning segment and an intracellular domain⁽¹⁸⁾. Most loss-of-function mutations contributing to OPPG, including homozygous mutations and compound heterozygotes, have been described in the highly conserved second β -propeller binding domain of the *LRP5* protein⁽¹⁹⁾.

Here we describe the first report of an atypical femoral fracture in a case of OPPG, explained by two novel missense *LRP5* mutations, and discuss the insights this provides. We further developed and validated *LRP5* protein models showing both heterozygous mutations destabilise their respective *LRP5* β -propellers; domains critical for protein and ligand binding.

CASE REPORTS

We report the case of a 40-year old white European man with OPPG. Congenital blindness was diagnosed at 6 months of age and osteoporosis aged 11 following his first 3 fractures (wrist aged 7, wrist aged 8, humerus aged 10). He suffered a low trauma fracture of the femur aged 16 leading to an abnormal gait due to unequal leg length. A bone biopsy at this time showed normal cortical thickness, normal mineralization, but severe trabecular osteoporosis. He sustained further fractures of the humerus aged 23 and then a unilateral atypical subtrochanteric femoral fracture aged 38 years when he slipped off a low curb whilst walking (Figure 1); he had received no anti-resorptive therapy before this time and had no prodromal pain. The fracture was managed by open reduction and internal fixation with a plate and screws (Supplemental figure S1).

Although a non-smoker, his alcohol consumption is high averaging 64 units per week; diet otherwise normal with unlimited exercise tolerance. He has no children. Extensive corneal scarring and iris hypoplasia in his right eye now limits his visual perception to light and dark. His left eye vision is severely limited by a persistent embryonic hyaloid amblyopia with convergent squint, nystagmus, raised intraocular pressure and cataract. On examination, he was obese (height 163cm, weight 105kg, BMI 39.5kg/cm², arm span 184cm), with both mild kyphosis and pectus carinatum, a mild leg length deficit, but no signs of joint disease. He had normal nails, sclera, hearing, dentition and cardiovascular and neurological examinations. Serum calcium, phosphate, alkaline phosphatase and fasting glucose were all normal, biochemistry showed no evidence of renal, inflammatory or endocrine disease. 25(OH) vitamin D level was 43 nmol/L; bone turnover markers were low [serum C-terminal collagen crosslinks (CTX) 0.02 µg/L and N-terminal procollagen (P1NP) 24 µg/L] before anti-resorptive treatment. Dual Energy X-ray Absorptiometry (DXA) scanning measured L1-L4 T-score -3.1, with femoral neck T-score -1.3 (Supplemental Figure S2). He received 9 months of oral alendronate with supplementary calcium and vitamin D after his atypical femoral fracture (aged 38), but this was discontinued after specialist review.

Our index case reported a 36-year old sister with very similar phenotype (Supplemental Figure S3): congenital blindness, early onset kyphosis, short stature, and four fractures to date, but no DNA sample was available. In contrast, their 57-year old mother, without ophthalmological disease, had only experienced a greenstick forearm fracture aged 8. After menarche at 13 and menopause at 53 years, she developed diet-controlled type II diabetes

mellitus. She consumes 30 units alcohol/week and smokes 30 cigarettes/day. Her DXA L1-L4 T-score was -2.7 and femoral neck T-score -1.7 (Supplemental Figure S4); FRAX assessment recommend lifestyle advice. There are no other cases of OPPG in her family. Our index case reported his 60-year old father (Supplemental Figure S3) to have no eye disease but, although fracture-free, he has developed a kyphosis with height loss. He has two further offspring by a second relationship; none show an OPPG phenotype. There was no history of consanguinity.

The family has never previously sought genetic testing as the mother had ascribed her offspring's phenotypes to her use of the antiemetic Debendox (dicucloamine-doxylamine-pyridoxine) during pregnancy, despite a series of epidemiological studies in the late 1970s finding no evidence of teratogenicity⁽²⁰⁻²²⁾.

METHODS

LRP5 Sequencing

Genomic DNA from blood was obtained using QIAamp DNA blood extraction kit (Qiagen, GmbH, Hilden, Germany) following manufacturer's indications. Written informed consent was obtained from the index case and his mother. Research complied with the World Medical Association Declaration of Helsinki-Ethical Principles for Medical Research Involving Human Subjects. We amplified by PCR and directly sequenced, using Sanger methodology, all 23 exons of human *LRP5* (RefSeq NM_002335) as well as the relevant exon-intron boundaries. Primers for PCR and sequencing were designed using Primer3 v4.0.0 software^(23;24) (Supplemental Table S1). Mutations found were confirmed on two independent PCR products.

In silico analysis of novel LRP5 mutations

Novel missense *LRP5* mutations were analysed using PolyPhen-2⁽²⁵⁾, SIFT⁽²⁶⁾, Pmut⁽²⁷⁾, PANTHER⁽²⁸⁾, Align-GVGD⁽²⁹⁾, and MutationTaster⁽³⁰⁾. Modelling of the four extracellular YWTD β -propeller/EGF-like domains of human *LRP5* (E1-E4) was undertaken using Modeller 9v12^(31;32) based upon highly similar crystal structure template domain-fragments of *LRP6* – E1-E2 (PDB ID: 4DG6) and E3-E4 (PDB ID: 4A0P). The *LRP5* E1-E2 and E3-E4 domains share ~76% and ~72% sequence identity (~86% sequence similarity in both cases) with the template *LRP6* sequences, respectively. The target-template alignment was based upon multiple sequence alignment among related *LRP5/6* protein sequences using ClustalX⁽³³⁾. Fifty models were generated using

Modeller 9v12, and the one with the lowest objective function score selected as the representative model in each case. The models were assessed using PROCHECK⁽³⁴⁾, WHAT IF^(35;36) and MetaMQAPII⁽³⁷⁾. Details about the protein model quality assessment are provided in the Supplementary Table S2. FoldX^(38;39) was used to estimate the free energy difference (stability change, expressed as $\Delta\Delta G$) upon mutagenesis from wild-type for the mutations.

RESULTS

PCR amplification and sequencing of *LRP5* revealed two novel missense mutations in our OPPG index case. We found in both the index case and his mother, a novel heterozygous c.2254C>T (p.R752W) mutation (Supplemental Figure S5A, S5B). Our OPPG index case also carried a second novel heterozygous missense mutation, c.235T>C (p.W79R) (Supplemental Figure S5C). *In silico* bioinformatic analysis suggested both mutations are deleterious (Supplemental Table S3). Both substituted amino acids are evolutionarily strictly conserved among *LRP5* orthologues and also among related *LRP4* and *LRP6* sequences⁽⁴⁰⁾ (Supplemental Figures S6 and S7).

In order to further assess potential impact of the mutations on the structure and function of *LRP5*, we created and validated three-dimensional (3-D) homology models for the four extracellular YWTD β -propeller/EGF-like domains (E1-E4) of *LRP5*. Mutation *W79R* is located in the second β -strand of blade 2 in the E1 domain, within the highly conserved “YWTD” signature motif, and is completely buried in the core of the β -propeller (Figure 2A). Protein stability calculations using FoldX on the 3-D model of *LRP5*-E1-E2 predict this mutation to severely destabilise structure (mean $\Delta\Delta G = \sim 5$ kcal/mol), based upon previously published studies^(39;41;42) ($\Delta\Delta G > 1.6$ kcal/mol infers severely reduced structural stability). The *R752W* mutation, located in blade 3 of E3 domain, near the interface with E4 (Figure 2B), is also predicted to destabilise structure, albeit to a lesser extent (mean $\Delta\Delta G = \sim 2$ kcal/mol).

DISCUSSION

To our knowledge, this is the first report of an atypical subtrochanteric femoral fracture occurring in a case of OPPG. Furthermore, we have identified two novel missense *LRP5* mutations explaining this compound

heterozygous case of OPPG, supported by 3-D protein models showing significant deleterious structural consequences to the LRP5 protein.

This AFF occurred within the context of relatively low bone turnover, impaired Wnt signalling due to predicted destabilisation of LRP5, and trabecular osteoporosis with normal mineralization on histology, which implicates impaired bone remodelling in the pathogenesis of AFF. Given our knowledge of LRP5 function, we hypothesize that this relatively low bone turnover state in adult OPPG may be chronic. Prolonged suppression of bone turnover with bisphosphonates, used to treat osteoporosis, has also been associated with AFF⁽¹⁾.

The underlying pathogenesis of AFF remains unclear. AFFs may represent a stress or insufficiency fracture, developing from accumulated cortical microcracks in response to repetitive loading, repair of which is hampered if bone remodelling is impaired⁽¹⁾. However, in contrast to stress fractures, AFFs usually start from the lateral (rather than medial) cortex, and fracture with a smooth transverse (rather than oblique) edge, suggestive of 'brittle bone', as seen in Figure 1. Femoral alignment, varus neck-shaft angle and shorter hip-axis length have been associated with AFF ^(43;44), with the initial cortical lesions occurring at the lateral femoral site under greatest tensile load ⁽⁴⁵⁾. Whether our index cases' mild leg-length deficit altered his lower limb biomechanics to increase tensile stress over the left lateral femur, predisposing to AFF, is unclear.

AFFs are thought to develop from a cortical defect. Earlier our index cases' bone biopsy had shown normal cortical thickness, but with trabecular osteoporosis. Clinical vertebral fractures have been reported more commonly in a handful of AFF cases⁽⁴⁶⁾. Whether a weakened trabecular scaffold within the femoral subtrochanteric region predisposes to cortical microcrack propagation is not known. Further lowering bone turnover by bisphosphonate therapy for OPPG, proposed by some in earlier case reports^(11;47), is inadvisable in our case, where arguably anabolic therapy would be the most appropriate therapeutic intervention.

AFFs have been reported in other rare bone diseases causing (i) impaired bone resorption due to osteoclast dysfunction with increased bone density *e.g.* pycnodysostosis⁽⁴⁸⁾, autosomal dominant type II osteopetrosis⁽⁴⁹⁻⁵¹⁾ (2 of 3 osteopetrosis cases occurred after high trauma accidents), and (ii) impaired mineralization due to hypophosphatasia⁽⁵²⁾. Until now AFF has never been reported in a genetic condition of primary osteoblast dysfunction.

We have identified two novel missense mutations in *LRP5*. Mutation *R752W*, within the E3 domain, is predicted to destabilise protein structure; however, the pathological effect is comparatively subtle, conferring only low BMD (no eye disease) in our postmenopausal mother who has other risk factors for low BMD, consistent with the mean $\Delta\Delta G$ just above the predicted destabilisation threshold (only ~ 0.4 kcal/mol above the 1.6 kcal/mol threshold). Interestingly, a *R752G* mutation has been described in familial exudative vitreoretinopathy⁽⁴⁰⁾; a rare retinal disorder with some cases showing reduced BMD⁽⁵³⁾. In addition to *R752W*, our index case has a second novel missense mutation; *W79R*, located within the core of the E1 β -propeller domain, is predicted to severely destabilise structure because of inappropriate burial of the polar/charged Arginine residue within a key conserved structural (YWTD) motif. Since we could not sequence either the father (normal vision) or sibling (with the same OPPG phenotype) we could not determine whether this second mutation is *de novo* or paternally inherited. Paternal inheritance is likely given the sibling's affection status and the father's height loss with kyphosis suggestive of osteoporosis.

LRP5 regulates bone formation through activation of osteoblastic Wnt/ β -catenin canonical signalling⁽⁵⁴⁾. Development of OPPG has been associated with homozygous and compound heterozygous mutations in *LRP5*^(2;6). No phenotypic differences have been reported between homozygotes and compound heterozygotes⁽⁶⁾. Interestingly, common polymorphisms in *LRP5* influence BMD in the general population⁽⁵⁵⁻⁶⁰⁾; the genetic architecture of individuals within the general population sustaining AFF is unknown.

It has been suggested that *LRP5* regulates bone mass through duodenal serotonin synthesis⁽⁶¹⁾, although subsequent analyses have not all supported this finding^(62;63). Furthermore, serotonin levels in high bone mass, due to activating *LRP5* mutations, have been reported to be normal⁽⁶⁴⁾; whether the same holds for OPPG remains to be determined.

In conclusion, we report the first case of atypical femoral fracture in a case of OPPG within the context of relatively low bone turnover, trabecular osteoporosis and impaired osteoblast function. The novel mutation *R752W* is associated with low BMD, but the combination of this mutation with another novel *W79R* mutation, causes a severe case of compound heterozygous OPPG, due to destabilisation of the β -propellers of the *LRP5* protein that are required for protein and ligand binding to regulate Wnt signalling and osteoblast function⁽⁶⁵⁾.

ACKNOWLEDGMENTS

We thank Ms Dilruba Kabir for her help with sequencing the *LRP5* gene. CLG is funded by Arthritis Research UK through a Clinician Scientist Fellowship (grant ref 20000). DCS acknowledges funding support from the Medical Research Council.

Author's roles: project conduct: SR, CLG, data collection: EM, GS, CLG, data analysis: NA, DCS, data interpretation: NA, DCS, CLG, drafting manuscript: NA, DCS, CLG, revising manuscript: NA, DCS, EM, GS, SR, CLG, approving final version of manuscript: NA, DCS, EM, GS, SR, CLG. NA takes responsibility for the integrity of the data analysis.

REFERENCES

- (1) Shane E, Burr D, Abrahamsen B, Adler RA, Brown TD, Cheung AM, et al. Atypical subtrochanteric and diaphyseal femoral fractures: second report of a task force of the american society for bone and mineral research. *J Bone Miner Res*. 2014 Jan;29(1):1-23.
- (2) Gong Y, Slee RB, Fukai N, Rawadi G, Roman-Roman S, Reginato AM, et al. LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell*. 2001 Nov 16;107(4):513-23.
- (3) Neuhauser G, Kaveggia EG, Opitz JM. Autosomal recessive syndrome of pseudogliomantous blindness, osteoporosis and mild mental retardation. *Clin Genet*. 1976 Mar;9(3):324-32.
- (4) Lee DH, Wenkert D, Whyte MP, Trese MT, Cruz OA. Congenital blindness and osteoporosis-pseudoglioma syndrome. *J AAPOS*. 2003 Feb;7(1):75-7.
- (5) Frontali M, Stomeo C, Dallapiccola B. Osteoporosis-pseudoglioma syndrome: report of three affected sibs and an overview. *Am J Med Genet*. 1985 Sep;22(1):35-47.
- (6) Ai M, Heeger S, Bartels CF, Schelling DK. Clinical and molecular findings in osteoporosis-pseudoglioma syndrome. *Am J Hum Genet*. 2005 Nov;77(5):741-53.
- (7) Cheung WM, Jin LY, Smith DK, Cheung PT, Kwan EY, Low L, et al. A family with osteoporosis pseudoglioma syndrome due to compound heterozygosity of two novel mutations in the LRP5 gene. *Bone*. 2006 Sep;39(3):470-6.
- (8) Marques-Pinheiro A, Levasseur R, Cormier C, Bonneau J, Boileau C, Varret M, et al. Novel LRP5 gene mutation in a patient with osteoporosis-pseudoglioma syndrome. *Joint Bone Spine*. 2010 Mar;77(2):151-3.
- (9) Narumi S, Numakura C, Shiihara T, Seiwa C, Nozaki Y, Yamagata T, et al. Various types of LRP5 mutations in four patients with osteoporosis-pseudoglioma syndrome: identification of a 7.2-kb microdeletion using oligonucleotide tiling microarray. *Am J Med Genet A*. 2010 Jan;152A(1):133-40.
- (10) Laine CM, Chung BD, Susic M, Prescott T, Semler O, Fiskerstrand T, et al. Novel mutations affecting LRP5 splicing in patients with osteoporosis-pseudoglioma syndrome (OPPG). *Eur J Hum Genet*. 2011 Aug;19(8):875-81.
- (11) Streeten EA, McBride D, Puffenberger E, Hoffman ME, Pollin TI, Donnelly P, et al. Osteoporosis-pseudoglioma syndrome: description of 9 new cases and beneficial response to bisphosphonates. *Bone*. 2008 Sep;43(3):584-90.
- (12) Levasseur R, LaCombe D, De Vernejoul MC. LRP5 mutations in osteoporosis-pseudoglioma syndrome and high-bone-mass disorders. *Joint Bone Spine*. 2005 May;72(3):207-14.
- (13) Saarinen A, Saukkonen T, Kivela T, Lahtinen U, Laine C, Somer M, et al. Low density lipoprotein receptor-related protein 5 (LRP5) mutations and osteoporosis, impaired glucose metabolism and hypercholesterolaemia. *Clin Endocrinol (Oxf)*. 2010 Apr;72(4):481-8.
- (14) Boyden LM, Mao J, Belsky J, Mitzner L, Farhi A, Mitnick MA, et al. High bone density due to a mutation in LDL-receptor-related protein 5. *N Engl J Med*. 2002 May 16;346(20):1513-21.
- (15) Little RD, Carulli JP, Del Mastro RG, Dupuis J, Osborne M, Folz C, et al. A mutation in the LDL receptor-related protein 5 gene results in the autosomal dominant high-bone-mass trait. *Am J Hum Genet*. 2002 Jan;70(1):11-9.

- (16) Van WL, Cleiren E, Gram J, Beals RK, Benichou O, Scopelliti D, et al. Six novel missense mutations in the LDL receptor-related protein 5 (LRP5) gene in different conditions with an increased bone density. *Am J Hum Genet.* 2003 Mar;72(3):763-71.
- (17) Ai M, Holmen SL, Van HW, Williams BO, Warman ML. Reduced affinity to and inhibition by DKK1 form a common mechanism by which high bone mass-associated missense mutations in LRP5 affect canonical Wnt signaling. *Mol Cell Biol.* 2005 Jun;25(12):4946-55.
- (18) Hey PJ, Twells RC, Phillips MS, Yusuke N, Brown SD, Kawaguchi Y, et al. Cloning of a novel member of the low-density lipoprotein receptor family. *Gene.* 1998 Aug 17;216(1):103-11.
- (19) Zhang Y, Wang Y, Li X, Zhang J, Mao J, Li Z, et al. The LRP5 high-bone-mass G171V mutation disrupts LRP5 interaction with Mesd. *Mol Cell Biol.* 2004 Jun;24(11):4677-84.
- (20) Fleming DM, Knox JD, Crombie DL. Debendox in early pregnancy and fetal malformation. *Br Med J (Clin Res Ed).* 1981 Jul 11;283(6284):99-101.
- (21) Milkovich L, van den Berg BJ. An evaluation of the teratogenicity of certain antinauseant drugs. *Am J Obstet Gynecol.* 1976 May 15;125(2):244-8.
- (22) Smithells RW, Sheppard S. Teratogenicity testing in humans: a method demonstrating safety of bendectin. *Teratology.* 1978 Feb;17(1):31-5.
- (23) Koressaar T, Remm M. Enhancements and modifications of primer design program Primer3. *Bioinformatics.* 2007 May 15;23(10):1289-91.
- (24) Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3--new capabilities and interfaces. *Nucleic Acids Res.* 2012 Aug;40(15):e115.
- (25) Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods.* 2010 Apr;7(4):248-9.
- (26) Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc.* 2009;4(7):1073-81.
- (27) Ferrer-Costa C, Gelpi JL, Zamakola L, Parraga I, de IC, X, Orozco M. PMUT: a web-based tool for the annotation of pathological mutations on proteins. *Bioinformatics.* 2005 Jul 15;21(14):3176-8.
- (28) Mi H, Muruganujan A, Thomas PD. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D377-D386.
- (29) Tavtigian SV, Deffenbaugh AM, Yin L, Judkins T, Scholl T, Samollow PB, et al. Comprehensive statistical study of 452 BRCA1 missense substitutions with classification of eight recurrent substitutions as neutral. *J Med Genet.* 2006 Apr;43(4):295-305.
- (30) Schwarz JM, Rodelsperger C, Schuelke M, Seelow D. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods.* 2010 Aug;7(8):575-6.
- (31) Sali A, Blundell TL. Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol.* 1993 Dec 5;234(3):779-815.
- (32) Fiser A, Sali A. Modeller: generation and refinement of homology-based protein structure models. *Methods Enzymol.* 2003;374:461-91.
- (33) Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 1997 Dec 15;25(24):4876-82.

- (34) Laskowski RA, MacArthur MW, Moss D.S., Thornton JM. Procheck - a Program to Check the Stereochemical Quality of Protein Structures. *J Appl Crystallogr.* 1993;26:283-91.
- (35) Vriend G. WHAT IF: a molecular modeling and drug design program. *J Mol Graph.* 1990 Mar;8(1):52-6, 29.
- (36) Vriend G, Sander C. Quality control of protein models: directional atomic contact analysis. *J Appl Crystallogr.* 1993 Jan 1;26:47-60.
- (37) Pawlowski M, Gajda MJ, Matlak R, Bujnicki JM. MetaMQAP: a meta-server for the quality assessment of protein models. *BMC Bioinformatics.* 2008;9:403.
- (38) Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field. *Nucleic Acids Res.* 2005 Jul 1;33(Web Server issue):W382-W388.
- (39) Guerois R, Nielsen JE, Serrano L. Predicting changes in the stability of proteins and protein complexes: a study of more than 1000 mutations. *J Mol Biol.* 2002 Jul 5;320(2):369-87.
- (40) Jiao X, Ventruto V, Trese MT, Shastry BS, Hejtmancik JF. Autosomal recessive familial exudative vitreoretinopathy is associated with mutations in LRP5. *Am J Hum Genet.* 2004 Nov;75(5):878-84.
- (41) Tokuriki N, Stricher F, Schymkowitz J, Serrano L, Tawfik DS. The stability effects of protein mutations appear to be universally distributed. *J Mol Biol.* 2007 Jun 22;369(5):1318-32.
- (42) Rakoczy EP, Kiel C, McKeone R, Stricher F, Serrano L. Analysis of disease-linked rhodopsin mutations based on structure, function, and protein stability calculations. *J Mol Biol.* 2011 Jan 14;405(2):584-606.
- (43) Saita Y, Ishijima M, Mogami A, Kubota M, Baba T, Kaketa T, et al. The fracture sites of atypical femoral fractures are associated with the weight-bearing lower limb alignment. *Bone.* 2014 Sep;66:105-10.
- (44) Taormina DP, Marcano AI, Karia R, Egol KA, Tejwani NC. Symptomatic atypical femoral fractures are related to underlying hip geometry. *Bone.* 2014 Jun;63:1-6.
- (45) Koh JS, Goh SK, Png MA, Ng AC, Howe TS. Distribution of atypical fractures and cortical stress lesions in the femur: implications on pathophysiology. *Singapore Med J.* 2011 Feb;52(2):77-80.
- (46) Giusti A, Hamdy NA, Dekkers OM, Ramautar SR, Dijkstra S, Papapoulos SE. Atypical fractures and bisphosphonate therapy: a cohort study of patients with femoral fracture with radiographic adjudication of fracture site and features. *Bone.* 2011 May 1;48(5):966-71.
- (47) Barros ER, Dias da Silva MR, Kunii IS, Lazaretti-Castro M. Three years follow-up of pamidronate therapy in two brothers with osteoporosis-pseudoglioma syndrome (OPPG) carrying an LRP5 mutation. *J Pediatr Endocrinol Metab.* 2008 Aug;21(8):811-8.
- (48) Yates CJ, Bartlett MJ, Ebeling PR. An atypical subtrochanteric femoral fracture from pycnodysostosis: a lesson from nature. *J Bone Miner Res.* 2011 Jun;26(6):1377-9.
- (49) Birmingham P, McHale KA. Case reports: treatment of subtrochanteric and ipsilateral femoral neck fractures in an adult with osteopetrosis. *Clin Orthop Relat Res.* 2008 Aug;466(8):2002-8.
- (50) Kumbaraci M, Karapinar L, Incesu M, Kaya A. Treatment of bilateral simultaneous subtrochanteric femur fractures with proximal femoral nail antirotation (PFNA) in a patient with osteopetrosis: case report and review of the literature. *J Orthop Sci.* 2013 May;18(3):486-9.
- (51) Sonohata M, Okubo T, Ono H, Mawatari M, Hotokebuchi T. Bipolar hip arthroplasty for subtrochanteric femoral nonunion in an adult with autosomal dominant osteopetrosis type II. *J Orthop Sci.* 2011 Sep;16(5):652-5.

- (52) Whyte MP. Atypical femoral fractures, bisphosphonates, and adult hypophosphatasia. *J Bone Miner Res.* 2009 Jun;24(6):1132-4.
- (53) Qin M, Hayashi H, Oshima K, Tahira T, Hayashi K, Kondo H. Complexity of the genotype-phenotype correlation in familial exudative vitreoretinopathy with mutations in the LRP5 and/or FZD4 genes. *Hum Mutat.* 2005 Aug;26(2):104-12.
- (54) Harada S, Rodan GA. Control of osteoblast function and regulation of bone mass. *Nature.* 2003 May 15;423(6937):349-55.
- (55) Baldock PA, Eisman JA. Genetic determinants of bone mass. *Curr Opin Rheumatol.* 2004 Jul;16(4):450-6.
- (56) Ferrari SL, Deutsch S, Choudhury U, Chevalley T, Bonjour JP, Dermitzakis ET, et al. Polymorphisms in the low-density lipoprotein receptor-related protein 5 (LRP5) gene are associated with variation in vertebral bone mass, vertebral bone size, and stature in whites. *Am J Hum Genet.* 2004 May;74(5):866-75.
- (57) Koay MA, Woon PY, Zhang Y, Miles LJ, Duncan EL, Ralston SH, et al. Influence of LRP5 polymorphisms on normal variation in BMD. *J Bone Miner Res.* 2004 Oct;19(10):1619-27.
- (58) Koay MA, Tobias JH, Leary SD, Steer CD, Vilarino-Guell C, Brown MA. The effect of LRP5 polymorphisms on bone mineral density is apparent in childhood. *Calcif Tissue Int.* 2007 Jul;81(1):1-9.
- (59) Saarinen A, Valimaki VV, Valimaki MJ, Loyttyniemi E, Auro K, Uusen P, et al. The A1330V polymorphism of the low-density lipoprotein receptor-related protein 5 gene (LRP5) associates with low peak bone mass in young healthy men. *Bone.* 2007 Apr;40(4):1006-12.
- (60) Sims AM, Shephard N, Carter K, Doan T, Dowling A, Duncan EL, et al. Genetic analyses in a sample of individuals with high or low BMD shows association with multiple Wnt pathway genes. *J Bone Miner Res.* 2008 Apr;23(4):499-506.
- (61) Yadav VK, Ryu JH, Suda N, Tanaka KF, Gingrich JA, Schutz G, et al. Lrp5 controls bone formation by inhibiting serotonin synthesis in the duodenum. *Cell.* 2008 Nov 28;135(5):825-37.
- (62) Cui Y, Niziolek PJ, MacDonald BT, Zylstra CR, Alenina N, Robinson DR, et al. Lrp5 functions in bone to regulate bone mass. *Nat Med.* 2011 Jun;17(6):684-91.
- (63) Chang MK, Kramer I, Keller H, Gooi JH, Collett C, Jenkins D, et al. Reversing LRP5-dependent osteoporosis and SOST deficiency-induced sclerosing bone disorders by altering WNT signaling activity. *J Bone Miner Res.* 2014 Jan;29(1):29-42.
- (64) Lee GS, Simpson C, Sun BH, Yao C, Foer D, Sullivan B, et al. Measurement of plasma, serum, and platelet serotonin in individuals with high bone mass and mutations in LRP5. *J Bone Miner Res.* 2014 Apr;29(4):976-81.
- (65) Balemans W, Van HW. The genetics of low-density lipoprotein receptor-related protein 5 in bone: a story of extremes. *Endocrinology.* 2007 Jun;148(6):2622-9.

FIGURE LEGENDS

Figure 1. X-ray showing a left atypical femoral fracture sustained by the index case aged 38. A subtrochanteric, transverse, non-comminuted fracture with medial spike is seen. Generalised increases in cortical thickness are seen in the diaphyses.

Figure 2. A) The 3-D model of LRP5 E1-E2 showing location of missense mutation W79R (sphere representation) in the second β -strand of blade 2 in the E1 domain. W79 is completely buried within the core of the β -propeller and the mutation is predicted to severely destabilise structure. A transparency setting was used for the surface representation to show the location of the buried residue on the secondary structure cartoon representation. Each of the blades that make up the six-bladed β -propeller structure in E1, and the four β -strands that make up blade 2 are labelled. **B)** Side-view of the 3-D model of LRP5 E3-E4 showing missense mutation R752W (sphere representation). R752 is located in blade 3 of the E3 domain, near the interface with E4.




Figure 1




Figure 2a




Figure 2b